

## Vaccination Schedule for Ewes and Lambs



### Lambs

1. Soremouth at docking/castration.
2. Tetanus antitoxin at docking/castration. If ewes were vaccinated with CDT, do not have to give this vaccination at docking/castration.
3. Overeating
  - a. Toxoid or bacterin at 5 weeks of age.
  - b. Toxoid or bacterin at 8 weeks of age (weaning).
  - c. Toxoid or bacterin at 11 weeks of age.

### Ewes

1. Vibriosis
  - a. 30 days before breeding.
  - b. Again, 3 to 4 months into gestation.
  - c. Booster every year 3 to 4 months into gestation.
2. Enzootic Abortion of Ewes (EAE)
  - a. 60 days before breeding.
  - b. 30 days later.
  - c. Booster every year 30 days before breeding.
3. CD/T
  - a. Give to lambs at 5, 8 and 11 weeks of age (can give CD/T, CD, or only D).
  - b. Booster to ewes every year 2 to 3 weeks before lambing begins.
  - c. If ewes were not vaccinated for CD/T as lambs, give 2 injections before first lambing, 6 and 2 weeks before lambing begins. Thereafter, give booster every year 2 to 3 weeks before lambing begins.
4. Caseous Lymphadenitis
  - a. 2 ml subcutaneous in spring (replacements).
  - b. Give booster 4 weeks later (replacements).
  - c. Give booster every spring.